arXiv:1007.1083v1 [math.AG] 7 Jul 2010

COBORDISM OF FLAG BUNDLES

AMALENDU KRISHNA

Abstract. Let G be a connected linear algebraic group over a field k of characteristic zero and let P be a parabolic subgroup of G containing a fixed maximal torus T. For a scheme X of finite type over k and a principal G-bundle $E \to X$, we describe the rational algebraic cobordism of the flag bundle $E/P \to X$ in terms of the cobordism groups of X and the classifying space BT. In particular, we obtain formulae for the algebraic cobordism groups of the various flag bundles associated to a vector bundle on a scheme. As a consequence, we describe the cobordism group of any principal bundle over a scheme. We also obtain similar formula for the higher Chow groups of flag bundles.

1. Introduction

Let k be a field of characteristic zero. In this paper, we shall consider only those schemes which are quasi-projective over k. Based on the construction of the motivic algebraic cobordism spectrum MGL by Voevodsky in the stable homotopy category of k, and the already known cobordism theory for complex manifolds [16], Levine and Morel [12] invented the algebraic cobordism theory $\Omega_*(-)$. The most important aspect of this theory is that $\Omega_*(-)$ is the universal oriented Borel-Moore homology theory in the category of k-schemes. In particular, it is the universal oriented cohomology theory in the category of smooth schemes over k.

As a consequence, many known theories, e.g., algebraic K-theory, Chow groups, can be directly obtained from the cobordism theory of Levine and Morel. This makes the question of describing the algebraic cobordism groups of various schemes interesting and important. Since this theory has been invented only some years ago, not many cases of computations of $\Omega_*(-)$ have been known. Levine and Morel showed that the coefficient ring $\Omega^*(k)$ is isomorphic to the known Lazard ring. They were also able to describe the algebraic cobordism of a projective bundle in terms of the cobordism group of the base scheme. The principal aim of this paper is to generalize this description to the case of arbitrary flag bundles. As a consequence, we also describe the cobordism groups of principal G-bundles over k-schemes.

In order to motivate our main results, we recall the following result, due to Borel and Leray, well known in algebraic topology and its analogue in algebraic geometry, due to Vistoli [19]. Assume $k = \mathbb{C}$ is the field of complex numbers and let G be a connected and reductive complex algebraic group. We fix a maximal torus T of G, a Borel subgroup B of G containing T and let W denote the Weyl group of G with respect to T. Let $\widehat{T}$ denote the character group of T and let $\mathrm{Sym}(\widehat{T})$ denote the symmetric algebra of $\widehat{T} \otimes \mathbb{Q}$ over $\mathbb{Q}$. Let X be a complex manifold and let $E \to X$ be a principal G-bundle. The reader can think of it as a $G(\mathbb{C})$-fiber bundle over $X(\mathbb{C})$. Since the principal bundles are represented by the maps to the classifying

---

2010 *Mathematics Subject Classification.* Primary 14C25; Secondary 19E15.

Key words and phrases. Algebraic cobordism, group actions, principal bundles.

spaces in topology, we immediately get the characteristic homomorphisms

$$\mathrm{Sym}(\widehat{T})^W \cong \mathrm{H}^*(BG, \mathbb{Q}) \xrightarrow{c_E} \mathrm{H}^*(X, \mathbb{Q}) \text{ and } \mathrm{Sym}(\widehat{T}) \xrightarrow{\alpha_E} \mathrm{H}^*(E/B, \mathbb{Q}).$$

The homomorphism α_E sends a character t of T to the first Chern class of the associated line bundle $E/B \overset{T,t}{\times} \mathbb{A}^1 \to E/B$. This induces a homomorphism of $\mathrm{Sym}(\widehat{T})$-algebras

$$\mathrm{H}^*(X, \mathbb{Q}) \otimes_{\mathrm{Sym}(\widehat{T})^W} \mathrm{Sym}(\widehat{T}) \xrightarrow{\lambda_X} \mathrm{H}^*(E/B, \mathbb{Q}) \tag{1.1}$$

and is an isomorphism.

Let X be now a scheme and let $\mathrm{CH}_*(X)$ denote the rational Chow group of algebraic cycles on X modulo the rational equivalence. Let $A^*(X)$ be the Fulton-MacPherson bivariant cohomology ring of X. Recall that this is a subring of the endomorphism ring of $\mathrm{CH}_*(X)$. In the same set up as above, Vistoli [19] showed that there are still the characteristic maps

$$\mathrm{Sym}(\widehat{T})^W \xrightarrow{c_E} A^*(X) \text{ and } \mathrm{Sym}(\widehat{T}) \xrightarrow{\alpha_E} A^*(E/B)$$

such that the induced map of $\mathrm{Sym}(\widehat{T})$-modules

$$\mathrm{CH}_*(X) \otimes_{\mathrm{Sym}(\widehat{T})^W} \mathrm{Sym}(\widehat{T}) \xrightarrow{\lambda_X} \mathrm{CH}_*(E/B) \tag{1.2}$$

is an isomorphism. This completely describes the Chow groups of the flag bundle in terms of the Chow group of the base.

In this paper, we study similar questions for the description of the higher Chow groups and more importantly, the algebraic cobordism groups of generalized flag bundles over any base scheme. The similar techniques can also be used to write down the description of the complex cobordism of flag bundles over complex manifolds. In case of the higher Chow groups, we obtain a more direct proof of the above formula using the localization sequence for these groups. In particular, this yields a different and simpler proof of Vistoli's theorem for the Chow groups. The proof in the case of the cobordism becomes much more complicated, mainly due to the absence of the *higher cobordism groups* at present. In this case, we adapt some of the arguments of [19] to the case of cobordism. We now state our main results.

Let G be a connected linear algebraic group and let T, B and W be a fixed split maximal torus, a Borel subgroup containing the maximal torus and the associated Weyl group respectively. We shall often denote this datum by the quadruplet (G, T, B, W). Let r denote the rank of T. Let P be a parabolic subgroup of G containing B and let W_P denote the Weyl group of the Levi subgroup of P with respect to T.

Let $S(G)$ and $C(G)$ denote the G-equivariant rational Chow ring and the algebraic cobordism ring of $\mathrm{Spec}(k)$ (see Section 3 below). Since we are interested in describing the higher Chow groups and cobordism groups with the rational coefficients, we make the convention that an abelian group A for us will actually mean $A \otimes_{\mathbb{Z}} \mathbb{Q}$. Furthermore, we shall write the higher Chow groups and the cobordism groups cohomologically in this paper in the sense that $\mathrm{CH}^i(X, n)$ and $\Omega^i(X)$ will mean the groups $\mathrm{CH}_{\dim(X)-i}(X, n)$ and $\Omega_{\dim(X)-i}(X)$ respectively. For any scheme X, we shall write the full Chow groups as

$$\mathrm{CH}^*(X) = \overset{\infty}{\underset{i=0}{\oplus}} \overset{\infty}{\underset{n=0}{\oplus}} \mathrm{CH}^i(X, n).$$

Let $p : E \to X$ be a principal G-bundle and let $\pi : E/P \to X$ be the flag bundle associated to the parabolic subgroup P. We show in Section 3 below that the algebraic cobordism groups $\Omega^*(X)$ and $\Omega^*(E/P)$ are modules over the rings $C(G)$ and $C(P)$ respectively. Moreover, it is known (*cf.* [10, Theorem 6.6]) that $C(G) = C(T)^W$ and $C(P) \cong C(T)^{W_P}$. The similar methods also show that the higher Chow groups $\mathrm{CH}^*(X, n)$ and $\mathrm{CH}^*(E/P, n)$ are modules over the rings $S(G) = S^W$ and $S(P) = S^{W_P}$ respectively. Now we have:

Theorem 1.1. *The natural map of $C(P)$-modules*

$$\lambda_X : \Omega^*(X) \otimes_{C(G)} C(P) \to \Omega^*(E/P) \tag{1.3}$$

is an isomorphism. Moreover, it is an isomorphism of rings if X is smooth.

Theorem 1.2. *The natural map of $S(P)$-modules*

$$\alpha_X : \mathrm{CH}^*(X) \otimes_{S(G)} S(P) \to \mathrm{CH}^*(E/P) \tag{1.4}$$

is an isomorphism. This is an isomorphism of rings if X is smooth.

As consequences of these results, we obtain the formulae (*cf.* Corollaries 6.6 and 8.2) for the cobordism and the higher Chow groups of principal bundles. We remark here that as we are working with the rational coefficients, the assumption about the maximal torus T being split is not a necessary one. One can reduce to this case by the transfer arguments.

We conclude the introduction with a brief outline of the contents of this paper. We recall the definitions and some important properties of the ordinary and the equivariant algebraic cobordism in the next section. We use these fundamental properties to construct our map λ_X in Section 3. We also deduce some functorial properties of this map with respect to morphisms between schemes. In section 4, we prove some algebraic results and use these together with some results of [10] to deduce our main result for algebraic cobordisms of trivial flag bundles over smooth schemes. In Section 5, we prove the surjectivity of λ_X using the localization sequence for the cobordism, the corresponding result for the trivial flag bundles and an induction argument. We prove Theorem 3.4 in Section 6 by first proving it for the trivial flag bundles, which uses the proof of the similar result for the Chow groups of trivial bundles in [19], and then using a filtration argument for general schemes. The final proof of Theorem 1.1 is given in Section 7, where we deduce this from the case of flag bundles associated to the Borel subgroups. The last section is devoted to the proof of Theorem 1.2 where the main tool is the long exact localization sequence for the higher Chow groups.

2. Recollection of ordinary and equivariant cobordism

In this section, we briefly recall the definitions and basic properties of the ordinary and equivariant algebraic cobordism.

2.1. **Algebraic cobordism.** Recall from [12] that for any scheme X and any $i \in \mathbb{Z}$, the algebraic cobordism group $\Omega^i(X)$ is given by the quotient of the $\mathbb{Q}$-vector space $Z^i(X)$ on the classes of projective morphisms $[Y \xrightarrow{f} X]$, where Y is a smooth scheme and f has relative codimension $i = \dim(X) - \dim(Y)$. This quotient is obtained by the relations in $Z^i(X)$ defined by certain axioms like the dimension axiom, section axiom and the formal group law. It was later shown by Levine and Pandharipande [14] that $\Omega^i(X)$ can also be described as the quotient of $Z^i(X)$ by the subspace generated by those cobordism cycles which are given by the *double point degeneration* relation. In particular, there is a natural surjection $Z^*(X) \twoheadrightarrow \Omega^*(X)$. It also follows that $\Omega^*(X)$ is a graded $\mathbb{Q}$-vector space, where the grading is given by the codimension of a cobordism cycle. Moreover, $\Omega^i(X) = 0$ for $i > \dim(X)$ and $\Omega^i(X)$ could be non-zero for any $-\infty < i \leq \dim(X)$. In fact, the exterior product on cobordism makes $\Omega^*(X)$ a graded ring for smooth X, which is a graded $\Omega^*(k)$-algebra. In general, $\Omega^*(X)$ is a graded $\Omega^*(k)$-module.

The following is the main result of Levine and Morel from which most of their other results on algebraic cobordism are deduced.

Theorem 2.1. *The functor $X \mapsto \Omega_*(X)$ is the universal Borel-Moore homology on the category of k-schemes. In other words, it is universal among the homology theories on this category which have functorial push-forward for projective morphism, pull-back for smooth morphism (any morphism of smooth schemes), Chern classes for line bundles, and which satisfy Projective bundle formula, homotopy invariance, the above dimension, section and formal group law axioms. Moreover, for a k-scheme X and closed subscheme Z of X of pure codimension p with open complement U, there is a localization exact sequence*

$$\Omega_*(Z) \to \Omega_*(X) \to \Omega_*(U) \to 0.$$

It was also shown in *loc. cit.* that the natural composite map

$$\Phi : \mathbb{L} \to \mathbb{L} \otimes_{\mathbb{Q}} \underline{\Omega}^*(k) \twoheadrightarrow \Omega^*(k)$$

$$a \mapsto [a]$$

is an isomorphism of commutative graded rings. Here, $\mathbb{L}$ is the Lazard ring which is a polynomial ring over $\mathbb{Q}$ on infinite but countably many variables and is given by the quotient of the polynomial ring $\mathbb{Q}[A_{ij} | (i,j) \in \mathbb{N}^2]$ by the relations, which uniquely define the universal formal group law $F_{\mathbb{L}}$ of rank one on $\mathbb{L}$. This formal group law is given by the power series

$$F_{\mathbb{L}}(u,v) = u + v + \sum_{i,j \geq 1} a_{ij} u^i v^j,$$

where a_{ij} is the equivalence class of A_{ij} in the ring $\mathbb{L}$. The Lazard ring is graded by putting the degree of a_{ij} to be $1-i-j$. In particular, one has $\mathbb{L}_0 = \mathbb{Q}, \mathbb{L}_{-1} = \mathbb{Q}a_{11}$ and $\mathbb{L}_i = 0$ for $i \geq 1$, that is, $\mathbb{L}$ is non-positively graded. We refer to *loc. cit.* for more properties of algebraic cobordism.

2.2. **Equivariant algebraic cobordism.**

Let (G, T, B, W) be the datum as above for a given connected linear algebraic group G over k. For a scheme X with a linear action of G, the equivariant algebraic cobordism of X was defined by Deshpande [4] when X is smooth and this was later defined and studied for all schemes in [10]. Since this is a new theory and since we shall have need for this here, albeit in a mild way, we briefly recall it.

For any integer $j \geq 0$, let V_j be an l-dimensional representation of G and let U_j be a G-invariant open subset of V_j such that the codimension of the complement $(V_j - U_j)$ in V_j is at least j and G acts freely on U_j such that the quotient U_j/G is a quasi-projective scheme. Such a pair (V_j, U_j) is called a *good* pair for the G-action corresponding to j. It is known that in our set up, good pairs always exist (*cf.* [5, Lemma 9]). Let X_G denote the mixed quotient $X \overset{G}{\times} U_j$ of the product $X \times U_j$ by the diagonal action of G, which is free.

Let X be a k-scheme of dimension d with a G-action. Fix $j \geq 0$ and let (V_j, U_j) be an l-dimensional good pair corresponding to j. Put

$$\Omega^G_i(X)_j = \frac{\Omega_{i+l-g}\left(X \overset{G}{\times} U_j\right)}{F_{d+l-g-j}\Omega_{i+l-g}\left(X \overset{G}{\times} U_j\right)}. \tag{2.1}$$

Here, $F_p\Omega_i(X)$ is the pth level of the Niveau filtration (*cf.* [10, Section 3]) which is roughly given by the subspace of $\Omega_i(X)$ generated by the images of $\Omega_i(Z) \to \Omega_i(X)$ under the push-forward map, where $Z \hookrightarrow X$ is a closed subscheme of dimension at most i. It is known that $\Omega_i^G(X)_j$ is independent of the choice of the good pair (V_j, U_j) and one defines

$$\Omega_i^G(X) := \varprojlim_j \Omega_i^G(X)_j. \tag{2.2}$$

The reader should note from the above definition that unlike the ordinary cobordism, the equivariant algebraic cobordism $\Omega_i^G(X)$ can be non-zero for any $i \in \mathbb{Z}$. We let

$$\Omega_*^G(X) = \bigoplus_{i\in\mathbb{Z}} \Omega_i^G(X)$$

and we let

$$\Omega^*_G(X) = \bigoplus_{i\in\mathbb{Z}} \Omega^i_G(X), \text{ where } \Omega^i_G(X) = \Omega^G_{\dim(X)-i}(X). \tag{2.3}$$

The equivariant cobordism satisfies all those properties which are listed in Theorem 2.1 for the ordinary algebraic cobordism. Since we shall need some of these properties, we state them below for the sake of completeness.

Theorem 2.2. *(cf.* [10, Theorems 5.1, 5.4]*) The equivariant algebraic cobordism satisfies the following properties.*
(i) Functoriality : The assignment $X \mapsto \Omega_*(X)$ is covariant for projective maps and contravariant for smooth maps of G-schemes. It is also contravariant for l.c.i. morphisms of G-schemes.
(ii) Homotopy : If $f : E \to X$ is a G-equivariant vector bundle, then $f^* : \Omega_*^G(X) \xrightarrow{\cong} \Omega_*^G(E)$.
(iii) Chern classes : For any G-equivariant vector bundle $E \xrightarrow{f} X$ of rank r, there are equivariant Chern class operators $c_l^G(E) : \Omega_*^G(X) \to \Omega_{*-l}^G(X)$ for $1 \le l \le r$ which have same functoriality properties as in the non-equivariant case.
(iv) Free action : If G acts freely on X with quotient Y, then $\Omega_*^G(X) \xrightarrow{\cong} \Omega_*(Y)$.
(v) Exterior Product : There is a natural product map

$$\Omega_i^G(X) \otimes_{\mathbb{Z}} \Omega_{i'}^G(X') \to \Omega_{i+i'}^G(X \times X').$$

In particular, $\Omega_*^G(k)$ is a graded algebra and $\Omega_*^G(X)$ is a graded $\Omega_*^G(k)$-module for every $X \in \mathcal{V}_G$.
(vi) Projection formula : For a projective map $f : X' \to X$ in $\mathcal{V}_G^S$, one has for $x \in \Omega_*^G(X)$ and $x' \in \Omega_*^G(X')$, the formula : $f_*(x' \cdot f^*(x)) = f_*(x') \cdot x$.
(vii) Localization sequence : For a G-invariant closed subscheme $Z \subset X$ with the complement U, there is an exact sequence

$$\Omega_*^G(Z) \to \Omega_*^G(X) \to \Omega_*^G(U) \to 0.$$

For a G-equivariant vector bundle E on X, we shall often denote the equivariant Chern class operators as $c_i^G(E) \cap -$. Note that these Chern classes behave like the Chern classes of the ordinary vector bundles on the ordinary cobordism of the mixed spaces defined before. In particular, if χ is a character of G (which is just a G-equivariant line bundle on spec(k)), the above exterior product is explicitly described as

$$\Omega^*_G(X) \otimes_{\mathbb{L}} \Omega^*(k) \to \Omega^*(X) \tag{2.4}$$

$$w \otimes c_1^G(L_\chi) \mapsto c_1^G\left(p^*(L_\chi)\right) \cap w$$

if $p : X \to \mathrm{Spec}(k)$ is the structure map. Here, L_χ is the line bundle associated to χ and $c_1^G(L_\chi)$ is identified with the element $c_1^G(L_\chi)(id)$ in the ring $C(G)$.

We shall denote $C(G) := \Omega^*(k)$ by $\Omega^*(BG)$ and call it as the cobordism ring of the *classifying space* of G. It is known from the universal property of the algebraic cobordism that for a complex linear algebraic group G, there is a natural map of rings

$$\rho_G : \Omega^*(BG) \to MU^*(BG), \tag{2.5}$$

where $MU^*(BG)$ is the rational complex cobordism group of the topological classifying space of $G(\mathbb{C})$. Moreover, this realization map is in fact an isomorphism (*cf.* [10, Theorem 6.8]). Thus, $\Omega^*(BG)$ is truly the cobordism ring of the classifying space of G.

If $H \subset G$ is a closed subgroup, then for any G-scheme X and a good pair (V_j, U_j), the G/H-fibration $X_H \to X_G$ induces a natural restriction map

$$r_H^G : \Omega_G^*(X) \to \Omega_H^*(X) \tag{2.6}$$

which in particular gives a natural $\mathbb{Q}$-algebra homomorphism $\Omega^*(BG) \to \Omega^*(BH)$. This restriction map in fact completely describes $\Omega^*(BG)$ in terms of $\Omega^*(BT)$ in the following way.

Theorem 2.3. *(cf.* [10, Theorem 6.6]*) The natural map $\Omega_G^*(X) \to \Omega_T^*(X)$ induces an isomorphism of $C(G)$-modules*

$$\Omega_G^*(X) \xrightarrow{\cong} \left(\Omega_T^*(X)\right)^W.$$

In particular, one has $C(G) \xrightarrow{\cong} C(T)^W$.

If T is a split torus of rank r and if $\{\chi_1, \cdots, \chi_r\}$ is a $\mathbb{Q}$-basis of $\widehat{T}$ with associated line bundles $\{L_{\chi_1}, \cdots, L_{\chi_r}\}$, then there is a natural ring isomorphism

$$\mathbb{L}[[x_1, \cdots, x_r]] \xrightarrow{\cong} C(T)$$

which maps x_i to the class of the first Chern class $c_1(L_{\chi_i})$ in $\Omega^*(BT)$. Similarly, there is an isomorphism $\mathbb{L}[[\gamma_1, \cdots, \gamma_n]] \xrightarrow{\cong} C(GL_n)$, where the image of γ_i is the ith Chern class of the canonical rank n vector bundle on BGL_n. Under the isomorphism $C(GL_n) \cong C(T)^W$ of Theorem 2.3, the image of γ_i is the ith elementary symmetric polynomial in the variables of $C(T)$.

We also recall here that there is a similar relation between the Chow rings of BG and BT (*cf.* [18], [5]), that is,

$$S(G) = \mathrm{CH}^*(BG) \xrightarrow{\cong} \mathrm{CH}^*(BT)^W = S(T)^W \tag{2.7}$$

and moreover

$$S(GL_n) \cong \mathbb{Q}[\gamma_1, \cdots, \gamma_n] \hookrightarrow \mathbb{Q}[x_1, \cdots, x_n] = S(T).$$

3. THE HOMOMORPHISM λ_X

In this section, we explain the homomorphism λ_X of Theorem 1.1 and then prove some functoriality properties of this map with respect to the maps between schemes. We consider the case of flag bundles associated to Borel subgroups of G, from which the general case can easily be deduced (*cf.* Section 7).

Let X be a scheme and let $p : E \to X$ be a principal G-bundle and let $\pi : E/B \to X$ be the flag bundle associated to the Borel subgroup B. Since E is a G-scheme where G acts freely, it follows from Theorem 2.2 that $\Omega^*(X) \cong \Omega_G^*(E)$

is a $C(G)$-modules. In the same way, $\Omega^*(E/T) \cong \Omega^*_T(E)$ is a $C(T)$-module (and hence a $C(G)$-module by restriction). On the other hand, $E/T \to E/B$ is a principal B^u-bundle, where B^u is the unipotent radical of B. By [3, XXII, 5.9.5], B^u has a finite filtration by normal subgroups whose successive quotients are the vector groups. A successive application of homotopy invariance now implies that $\Omega^*(E/B) \xrightarrow{\cong} \Omega^*(E/T)$. Hence, $\Omega^*(E/B)$ is a $C(T)$-module. Thus, $\Omega^*(X)$ and $\Omega^*(E/B)$ are naturally $C(G)$ and $C(T)$-modules respectively, which defines the map λ_X as $\lambda_X(w \otimes x) = x \cdot \pi^*(w)$.

Recall that $C(T)$ is the power series over $\mathbb{L}$ in the first Chern classes of the line bundles associated to the characters of T and Theorem 2.3 implies that $C(G)$ is also generated by the first Chern classes of the W-invariant characters inside $C(T)$.

Using the description of these module structures in (2.4), we see that the map λ_X is given by

$$\begin{aligned} \lambda_X : \Omega^*(X) \otimes_{C(G)} C(T) &\to \Omega^*(E/B) \\ w \otimes c_1(\chi) &\mapsto c_1(\chi) \cap \pi^*(w) \end{aligned} \tag{3.1}$$

for a character χ of T. It is easy to see from this that this is an $\mathbb{L}$-algebra homomorphism if X is smooth.

Remark 3.1. For readers who are little bit familiar with the language of quotient stacks and know that the G-equivariant line bundles on a G-scheme X are same as ordinary line bundles on the quotient stack $[X/G]$, we can explain the above in this set up as follows. The principal G-bundle $E \to X$ uniquely gives rise to the following commutative diagram of morphisms.

$$\begin{tikzcd} E/B \arrow[r, "\pi"] \arrow[d, "q"'] & X \arrow[d, "\overline{p}"] \\ BT \arrow[r] & BG, \end{tikzcd} \tag{3.2}$$

where BG is the quotient stack $[k/G]$. The map λ_X is then given as $\lambda_X\left(w \otimes c_1(\chi)\right) = c_1(q^*(L_\chi)) \cap \pi^*(w)$.

In particular, if $E = G \times X \to X$ is a trivial principal bundle, then the map $X \to BG$ canonically factors through the structure map $X \to \operatorname{Spec}(k) \to [k/G] = BG$. Hence, the map λ_X in this case is given by

$$\Omega^*(X) \otimes_{\mathbb{L}} \left(\mathbb{L} \otimes_{C(G)} C(T)\right) \to \Omega^*(E/B). \tag{3.3}$$

Here, the left term is identified as $\Omega^*(X) \otimes_{\mathbb{L}} \Omega^*(G/B)$ by [10, Theorem 7.6] and $\Omega^*(X) \otimes_{\mathbb{L}} \Omega^*(G/B) \xrightarrow{\lambda_X} \Omega^*(E/B)$ is simply the exterior product map.

To prove certain functoriality properties of λ_X, we need the following elementary result on the equivariant cobordism.

Lemma 3.2. *Let G be a linear algebraic group over k and let $f : Y \to X$ and $g : Z \to X$ be projective and smooth morphisms of G-schemes respectively. Then the maps $f_* : \Omega^*_G(Y) \to \Omega^*_G(X)$ and $g^* : \Omega^*_G(X) \to \Omega^*_G(Z)$ are $C(G)$-linear.*

Proof. We only give a sketch for the $C(G)$-linearity of f_*. The assertion about g^* is similar and much simpler. Note that the $C(G)$-module structure on $\Omega^*_G(X)$ is given by the exterior product (*cf.* Theorem 2.2). It suffices to show that

$$f_*(x \cdot w) = x \cdot f_*(w)$$

when x and w are generators of the corresponding cobordism groups $\Omega^*(U_j/G)$ and $\Omega^*((Y \times U_j)/G)$, where (V_j, U_j) is any given good pair for G-action. So let

$W_1 \xrightarrow{s_1} U_j/G$ and $W_2 \xrightarrow{s_2} Y \overset{G}{\times} U_j$ be projective morphisms from smooth and connected schemes, representing the cobordism classes x and w respectively. Let $\widetilde{W_1}$ and $\widetilde{W_2}$ be the pull-backs of W_1 and W_2 to U_j and $Y \times U_j$ respectively. By the definition of the push-forward and exterior product, we have

$$\begin{aligned} f_*(x \cdot w) &= f_*\left(\left[\widetilde{W_1} \overset{G}{\times} \widetilde{W_2} \to Y \overset{G}{\times} (U_j \times U_j)\right]\right) \\ &= \left[\widetilde{W_1} \overset{G}{\times} \widetilde{W_2} \to X \overset{G}{\times} (U_j \times U_j)\right] \end{aligned}$$

and the last term is same as the class of $x \cdot f_*(w)$ in $\Omega^*(X_G)$ which can be taken as $X \overset{G}{\times} (U_j \times U_j)$ because $(V_j \times V_j, U_j \times U_j)$ is also a good pair for the G-action. □

Lemma 3.3. *Let $f : Y \to X$ and $g : Z \to X$ be respectively, the projective and the smooth morphisms. Let $p : E \to X$ be a principal G-bundle and let E_Y and E_Z denote its pull-backs to Y and Z respectively. Consider the following Cartesian diagrams of flag bundles.*

(3.4)
$$\begin{tikzcd} E_Y/B \arrow[r, "\overline{f}"] \arrow[d, "\pi_Y"'] & E/B \arrow[d, "\pi"] \\ Y \arrow[r, "f"'] & X \end{tikzcd} \qquad \begin{tikzcd} E_Z/B \arrow[r, "\overline{g}"] \arrow[d, "\pi_Z"'] & E/B \arrow[d, "\pi"] \\ Z \arrow[r, "g"'] & X. \end{tikzcd}$$

Then the diagrams

(3.5)
$$\begin{tikzcd} \Omega^*(Y) \otimes_{C(G)} C(T) \arrow[r, "\lambda_Y"] \arrow[d, "f_* \otimes \mathrm{id}"'] & \Omega^*(E_Y/B) \arrow[d, "\overline{f}_*"] \\ \Omega^*(X) \otimes_{C(G)} C(T) \arrow[r, "\lambda_X"'] & \Omega^*(E/B) \end{tikzcd}$$

(3.6)
$$\begin{tikzcd} \Omega^*(X) \otimes_{C(G)} C(T) \arrow[r, "\lambda_X"] \arrow[d, "g^* \otimes \mathrm{id}"'] & \Omega^*(E/B) \arrow[d, "\overline{g}^*"] \\ \Omega^*(Z) \otimes_{C(G)} C(T) \arrow[r, "\lambda_Z"'] & \Omega^*(E_Z/B) \end{tikzcd}$$

are commutative.

Proof. To show the commutativity of the first square, we have

$$\begin{aligned} \overline{f}_* \circ \lambda_Y(w \otimes x) &= \overline{f}_*(x \cdot \pi_Y^*(w)) \\ &= x \cdot \overline{f}_*(\pi_Y^*(w)) \qquad \text{(By Lemma 3.2)} \\ &= x \cdot \pi^*(f_*(w)) \\ &= \lambda_X(f_*(w) \otimes x), \end{aligned}$$

where the third equality follows from the fact that the first square in (3.4) is Cartesian with π smooth and f projective. The proof of the commutativity of the second square is similar. □

Let (G, T, B, W) be as above where G is a connected linear algebraic group. Let G^u denote the unipotent radical of G and let L denote the corresponding quotient as a reductive group. Then any principal G-bundle $E \to X$ canonically gives a principal L-bundle $E_L = E/G^u \to X$. Moreover, as the Borel subgroup B contains G^u, we see that $E/B \xrightarrow{\cong} E_L/B_L$, where B_L is the image of B which is a Borel subgroup of the reductive group L. Since $C(G) \cong C(L)$, as follows from the Levi decomposition and the homotopy invariance, we conclude that it is enough to consider the case when G is reductive in order to prove our main results. Hence for the rest of this paper, G will always denote a connected reductive group. We shall deduce Theorem 1.1 from the following result for the algebraic cobordism of the flag bundles associated to the Borel subgroup B.

Theorem 3.4. *Let $p : E \to X$ be a principal G-bundle and let $\pi : E/B \to X$ be the flag bundle associated to the Borel subgroup B. The natural map of $C(T)$-modules*

$$\lambda_X : \Omega^*(X) \otimes_{C(G)} C(T) \to \Omega^*(E/B) \tag{3.7}$$

is an isomorphism. Moreover, it is an isomorphism of rings if X is smooth.

4. Some algebraic reductions

Let (G, T, B, W) be as above and let T be a split torus of rank r. This rank will be fixed throughout. We fix a basis $\{\chi_1, \cdots, \chi_r\}$ of $\widehat{T}$ and let $S = \mathrm{Sym}(\widehat{T}) = \mathbb{Q}[x_1, \cdots, x_r]$ be the polynomial algebra in the first Chern classes of the line bundles associated to the characters $\{\chi_i, \cdots, \chi_r\}$. Let $S^W \subset S$ be the subalgebra generated by the homogeneous polynomials which are invariant under the action of W. This gives us a square of ring inclusions

$$\begin{tikzcd}
(\mathbb{L}[x_1, \cdots, x_r])^W \arrow[r] \arrow[d] & \mathbb{L}[x_1, \cdots, x_r] \arrow[d] \\
C(G) \arrow[r] & C(T),
\end{tikzcd} \tag{4.1}$$

which is Cartesian and where $C(G)$ has been identified with $C(T)^W$. We shall write $\mathbb{L}[x_1, \cdots, x_r]$ simply as $S_{\mathbb{L}}$. Note that $S_{\mathbb{L}}$ and $S^W_{\mathbb{L}}$ are canonically isomorphic to $\mathbb{L} \otimes_{\mathbb{Q}} S$ and $\mathbb{L} \otimes_{\mathbb{Q}} S^W$ as $\mathbb{L}$-algebras. It is also known that $S^W_{\mathbb{L}}$ is a polynomial algebra over $\mathbb{L}$ of rank r. We shall denote the homogeneous generators of this subalgebra by $\{\sigma_1, \cdots, \sigma_r\}$.

Let I be the ideal of S generated by the homogeneous elements of positive degree which are invariant under W and let Λ denote the ring S/I. Then we see that $\Lambda_{\mathbb{L}} = \mathbb{L} \otimes_{\mathbb{Q}} \Lambda$ is canonically isomorphic to the $\mathbb{L}$-algebra $\mathbb{L}[x_1, \cdots, x_r]/I$. We recall the following result from [19, Lemma 1.2].

Lemma 4.1. *The graded $\mathbb{Q}$-algebra Λ is finite. If N is the maximal integer for which $\Lambda_N \neq 0$, then $N = \dim(G/B)$. Moreover, the $\mathbb{Q}$-vector space Λ_N is one-dimensional, and if d is an integer, the homomorphism*

$$\Lambda_d \otimes \Lambda_{N-d} \to \Lambda_N$$

given by the multiplication in Λ is a perfect pairing of finite-dimensional $\mathbb{Q}$-vector spaces.

Lemma 4.2. *Let A be a commutative ring and let I be an ideal of A. Let J be a finitely generated ideal of A. Then for any A-module M, the natural maps of $\widehat{A}$-modules*

$$J\widehat{M} \to \widehat{JM}, \quad \frac{\widehat{M}}{J\widehat{M}} \to \widehat{\left(\frac{M}{JM}\right)}$$

are isomorphisms, where $\widehat{M}$ denotes the I-adic completion of M.

Proof. Consider the exact sequence

$$0 \to JM \to M \to \frac{M}{JM} \to 0. \tag{4.2}$$

Since the topology on M is given by the descending chain $M \supset IM \supset I^2M \supset \cdots$ of submodules, it follows from [15, Theorem 8.1] that

$$0 \to \widehat{(JM)} \to \widehat{M} \to \widehat{\left(\frac{M}{JM}\right)} \to 0 \tag{4.3}$$

is exact. Thus, we only need to show the first isomorphism to prove the lemma.

Suppose $J = \sum\limits_{i=1}^{n} a_iA$ and define

$$\phi : M^n \to M$$

$$\phi(m_1, \cdots, m_n) = \sum_{i=1}^{n} a_i m_i.$$

This makes the sequence

$$M^r \to M \to \frac{M}{JM} \to 0 \tag{4.4}$$

exact. It again follows from [15, Theorem 8.1] that

$$\widehat{M^r} \xrightarrow{\widehat{\phi}} \widehat{M} \to \widehat{\left(\frac{M}{JM}\right)} \to 0 \tag{4.5}$$

is exact. On the other hand, $\widehat{\phi}$ is again given by $\widehat{\phi}(\widehat{m_1}, \cdots, \widehat{m_n}) = \sum\limits_{i=1}^{n} a_i\widehat{m_i}$. In other words, Image$(\widehat{\phi}) = J\widehat{M}$. The first isomorphism now follows from this and (4.3). This proves the lemma. □

Corollary 4.3. *The natural homomorphisms of rings*

$$\mathbb{L} \otimes \Lambda \xrightarrow{\cong} \frac{\mathbb{L}[x_1, \cdots, x_r]}{I} \to \frac{C(T)}{IC(T)} \to C(T) \otimes_{C(G)} \mathbb{L}$$

are isomorphisms.

Proof. We first observe that the cobordism ring $C(T)$ is the inverse limit of the cobordism rings of the form $\left(\Omega^*(BT)\right)_{j \geq 0}$ on each of which the Weyl group acts. In particular, the action of W on $C(T)$ is induced by its action on the polynomial ring $\mathbb{L}[x_1, \cdots, x_r]$ and $C(T)^W$ is the inverse limit of the W-invariants in the inverse system $\left(\Omega^*(BT)\right)_{j \geq 0}$. Thus we see that we can write $S^W_{\mathbb{L}} = \mathbb{L}[\sigma_1, \cdots, \sigma_r] \hookrightarrow \mathbb{L}[x_1, \cdots, x_r]$ and $C(G) = C(T)^W$ is the subring of the power series ring $\mathbb{L}[[x_1, \cdots, x_r]]$ generated by the homogeneous polynomials $\{\sigma_1, \cdots, \sigma_r\}$. Moreover, the ideal I in

$C(T)$ is the extension of the ideal $(\sigma_1, \cdots, \sigma_r)$ of $S^W = \mathbb{Q}[\sigma_1, \cdots, \sigma_r]$ which we also denote by I. Let $\mathfrak{m}$ denote the ideal $(x_1, \cdots, x_r)$ of $S_{\mathbb{L}}$. Now we have

$$\begin{aligned} C(T) \otimes_{C(G)} \mathbb{L} &\cong C(T) \otimes_{C(G)} \left(\frac{C(G)}{(\sigma_1, \cdots, \sigma_r)}\right) \\ &\cong \frac{C(T)}{IC(T)} \\ &\cong \frac{\widehat{(S_{\mathbb{L}})_{\mathfrak{m}}}}{I\widehat{(S_{\mathbb{L}})_{\mathfrak{m}}}} \\ &\cong \widehat{\left(\frac{S_{\mathbb{L}}}{IS_{\mathbb{L}}}\right)_{\mathfrak{m}}} \qquad \text{(By Lemma 4.2).} \end{aligned}$$

On the other hand,

$$\frac{S_{\mathbb{L}}}{IS_{\mathbb{L}}} \cong \mathbb{L} \otimes_{\mathbb{Q}} \left(\frac{\mathbb{Q}[x_1, \cdots, x_r]}{I}\right) \cong \prod_{j=1}^{s} \mathbb{L} \otimes_{\mathbb{Q}} A_j, \tag{4.6}$$

where each A_j is an artinian local ring which is finite over $\mathbb{Q}$. In particular, the ideal $\mathfrak{m}$ is nilpotent in each of the factor $\mathbb{L} \otimes A_j$ and hence the last term in (4.6) is complete with respect to $\mathfrak{m}$. We conclude that $\frac{S_{\mathbb{L}}}{IS_{\mathbb{L}}}$ is complete in the $\mathfrak{m}$-adic topology. In particular, we obtain

$$C(T) \otimes_{C(G)} \mathbb{L} \cong \widehat{\left(\frac{S_{\mathbb{L}}}{IS_{\mathbb{L}}}\right)_{\mathfrak{m}}} \cong \frac{S_{\mathbb{L}}}{IS_{\mathbb{L}}}$$

and this completes the proof. □

Corollary 4.4. *Let $X \times G/B \xrightarrow{\pi} X$ be the trivial flag bundle. Then the map λ_X is given by*

$$\Omega^*(X) \otimes_{\mathbb{Q}} \Lambda \xrightarrow{\cong} \Omega^*(X) \otimes_{\mathbb{L}} \Lambda_{\mathbb{L}} \xrightarrow{\lambda_X} \Omega^*(E/B)$$

which is an $\mathbb{L}$-algebra isomorphism if X is smooth.

Proof. The first assertion of the corollary follows directly from (3.3) and Corollary 4.3. If X is smooth, this map is an $\mathbb{L}$-algebra homomorphism because so are the maps in (3.1) and Corollary 4.3. Moreover, it is an isomorphism by [10, Lemma 6.5] and [13, Theorem 3.1]. □

5. Surjectivity of λ_X

We now let $p : E \to X$ be a an arbitrary principal G-bundle and let $\pi : E/B \to X$ be the associated flag bundle. Using the above inclusions of the polynomial rings inside the power series rings, we get natural homomorphisms

$$\begin{tikzcd} \Omega^*(X) \otimes_{S_{\mathbb{L}}^W} S_{\mathbb{L}} \arrow[d, "\delta_X"'] \arrow[dr, "\phi_X"] & \\ \Omega^*(X) \otimes_{C(G)} C(T) \arrow[r, "\lambda_X"'] & \Omega^*(E/B) \end{tikzcd} \tag{5.1}$$

of $S_{\mathbb{L}}$-modules, which are also $S_{\mathbb{L}}$-algebra homomorphisms if X is smooth.

As a first step towards proving Theorem 3.4, we show in this section that the map λ_X is surjective. In fact, the proof that follows will show that the map ϕ_X is surjective. It will eventually turn out that both the maps ϕ_X and λ_X are isomorphisms. We begin with the following elementary property of principal bundles and the local property of algebraic cobordism.

Lemma 5.1. *Let $p : E \to X$ be a principal G-bundle and let $\pi : E/B \to X$ be the associated flag bundle. Then the G-action $G \times E \xrightarrow{\mu} E$ induces a commutative diagram*

(5.2)
$$\begin{tikzcd}
G/B \times E/B \arrow[r, "\mu'"] \arrow[d, "\pi'"'] & E/B \arrow[d, "\pi"] \\
E/B \arrow[r, "\pi"'] & X
\end{tikzcd}$$

which is Cartesian and where π' is the projection to the second factor.

Proof. Since $E \xrightarrow{p} X$ is a principal G-bundle quotient of quasi-projective schemes, the action map $G \times E \xrightarrow{\mu} E$ induces a commutative diagram

(5.3)
$$\begin{tikzcd}
G \times E \arrow[r, "\mu"] \arrow[d, "p'"'] & E \arrow[d, "p"] \\
E \arrow[r, "p"'] & X
\end{tikzcd}$$

which is Cartesian and where p' is the projection to the second factor by the general properties of principal bundles (*cf.* [7, 0.10]).

Now, the map μ descends to a map $G \times E/B \to E/B$. Moreover, this map is B-equivariant where B acts trivially on E/B and by left multiplication on G. Taking the quotients, we get a canonical map $G/B \times E/B \xrightarrow{\mu'} E/B$ making the diagram (5.2) commute. It is now an easy exercise to check from (5.3) that this diagram is Cartesian too. □

Lemma 5.2. *Let $f : X' \to X$ be a finite and étale morphism of smooth and connected schemes. Then there exists an open subscheme $U \overset{j}{\hookrightarrow} X$ such that for the map $g = f|_{U'} : U' = f^{-1}(U) \to U$, one has $g_*(1) = [k(X') : k(X)]$.*

Proof. Let η denote the generic point of X and consider the Cartesian diagram

(5.4)
$$\begin{tikzcd}
X'_\eta \arrow[r, "i'"] \arrow[d, "h"'] & U' \arrow[r, "j'"] \arrow[d, "g"] & X' \arrow[d, "f"] \\
\eta \arrow[r, "i"'] & U \arrow[r, "j"'] & X,
\end{tikzcd}$$

where U is any open subscheme of X. Since X' is connected, we see that $X_\eta = \mathrm{Spec}(k(X'))$. Put $p = j \circ i$, $p' = j' \circ i'$ and $d = [k(X') : k(X)]$. It follows from [11, Lemma 4.7] that

$$p^* \circ f_*(1) = h_* \circ p'^*(1) = h_*(1) = d.$$

Since the algebraic cobordism is generically constant by [11, Lemma 13.3, Corollary 13.4], there exists an open subscheme $U \overset{j}{\hookrightarrow} X$ such that $j^* \circ f_*(1) = d$ in $\Omega^*(U)$. This in turn implies that

$$g_*(1) = g_* \circ j'^*(1) = j^* \circ f_*(1) = d$$

and this proves the lemma. □

Corollary 5.3. *Let* $f : X' \to X$ *be a finite and étale morphism of smooth and connected schemes and consider the diagram* (3.4). *Then there exists an open subscheme* $U \overset{j}{\hookrightarrow} X$ *such that for* $g = f|_{U'} : U' = f^{-1}(U) \to U$, *one has a commutative diagram*

(5.5)
$$\begin{tikzcd}
\Omega^*(U) \otimes_{C(G)} C(T) \arrow[r, "g^*"] \arrow[d, "\lambda_U"'] & \Omega^*(U') \otimes_{C(G)} C(T) \arrow[r, "g_*"] \arrow[d, "\lambda_{U'}"'] & \Omega^*(U) \otimes_{C(G)} C(T) \arrow[d, "\lambda_U"'] \\
\Omega^*(E_U/B) \arrow[r, "\overline{g}^*"'] & \Omega^*(E_{U'}/B) \arrow[r, "\overline{g}_*"'] & \Omega^*(E_U/B)
\end{tikzcd}$$

such that the horizontal composite maps are multiplication by $[k(X') : k(X)]$.

Proof. We choose $U \overset{j}{\hookrightarrow} X$ as in Lemma 5.2. The commutativity of the diagram follows from Lemma 3.3. Moreover, as f is finite and étale of degree d, it follows that $\overline{f}$ is also a morphism of the same type. We claim that $\overline{g}_*(1) = k(X') : k(X)]$. To see this, we evaluate the required term as

$$\overline{g}_*(1) = \overline{g}_* \circ \pi_{U'}^*(1) = \pi_U^* \circ g_*(1) = [k(X') : k(X)],$$

where the second equality follows from Lemma 5.2 and this proves the claim. The corollary now follows from the projection formula. □

Proposition 5.4. *The map* λ_X *is surjective for any scheme* X.

Proof. We shall prove this by induction on the dimension of X. We can assume that X is reduced. If X is zero-dimensional, it is of the form $X = \mathrm{Spec}(K)$, where K is a finite product of finite field extensions of k. We prove the case when $X = \mathrm{Spec}(k)$. The same proof applies for any finite extension of k. Now, there is a finite extension $k \hookrightarrow l$ such that Y_l is of the form G_l/B_l. Hence the result holds for $X = \mathrm{Spec}(l)$ by [10, Theorem 7.6]. The case of $\mathrm{Spec}(k)$ now follows from Corollary 5.3.

If the map π is of the form $X \times G/B \xrightarrow{\pi} X$ with X smooth, the maps λ_X and ϕ_X are in fact isomorphisms by (3.3), Corollary 4.4, [10, Lemma 6.5] and [13, Theorem 3.1]. In the general case, we can find an étale cover $X' \xrightarrow{f} X$ such that the base change $E/B \times_X X' \xrightarrow{\pi'} X'$ is the trivial flag bundle $X' \times G/B \to X'$. We can now find a smooth and dense open subset $U \overset{j}{\hookrightarrow} X$ such that the map $U' = f^{-1}(U) \to U$ is finite and étale. Moreover, the flag bundle is still trivial on U'. Since U is a disjoint union of smooth and connected schemes, the surjectivity of λ_U follows from the case of the trivial bundle shown above and Corollary 5.3.

We now let $Z = X - U$ be the complement of U in X with the reduced closed subscheme structure and consider the diagram

(5.6)
$$\begin{tikzcd}
\Omega^*(Z) \otimes_{C(G)} C(T) \arrow[r] \arrow[d, "\lambda_Z"'] & \Omega^*(X) \otimes_{C(G)} C(T) \arrow[r] \arrow[d, "\lambda_X"'] & \Omega^*(U) \otimes_{C(G)} C(T) \arrow[r] \arrow[d, "\lambda_U"] & 0 \\
\Omega^*(E_Z/B) \arrow[r] & \Omega^*(E/B) \arrow[r] & \Omega^*(E_U/B) \arrow[r] & 0
\end{tikzcd}$$

which is commutative by Lemma 3.3 and whose rows are exact by Theorem 2.1. Since U is open and dense, Z is a closed subscheme of dimension which is strictly smaller than that of X. Hence the map λ_Z is surjective by induction. We have shown above that λ_U is surjective. Hence the map λ_X is surjective too. □

Remark 5.5. Since the maps

$$\Omega^*(X) \otimes_{S^W_{\mathbb{L}}} S_{\mathbb{L}} \xrightarrow{\delta_X} \Omega^*(X) \otimes_{C(G)} C(T) \xrightarrow{\lambda_X} \Omega^*(E/B)$$

are isomorphisms for the trivial bundle $G/B \times X \xrightarrow{\pi} X$ for X smooth by Corollary 4.4, exactly the same proof as for Proposition 5.4 shows that the maps δ_X and ϕ_X are also surjective for any scheme X.

6. Proof of Theorem 3.4

Recall from Section 4 that $\Lambda = S/I = \Lambda_0 \oplus \Lambda_1 \oplus \cdots \oplus \Lambda_N$ is the graded quotient of $S = \mathbb{Q}[x_1, \cdots, x_r]$ by the ideal I which is generated by the homogeneous polynomials of positive degree which are invariant under W. It is clear that Λ_0 is one-dimensional over $\mathbb{Q}$ generated by the unit element of the ring. It also follows from Lemma 4.1 that Λ_N is an one-dimensional $\mathbb{Q}$-vector space. We fix these two generators and denote them by $\rho_0 = 1$ and ρ_N respectively. Let $p_0 = 1$ and p_N be their homogeneous lifts in S_0 and S_N respectively. For any scheme X, we consider the map

$$\psi : \Omega^*(X) \to \Omega^*(X) \tag{6.1}$$

$$\psi(x) = \pi_* \circ \phi_X(x \otimes p_N) = \pi_* \circ (c_1(p_N) \cap \pi^*(x))$$

where ϕ_X is the homomorphism in (5.1). We need the following property of this map.

Lemma 6.1. *Let $Z \overset{i}{\hookrightarrow} X$ be a closed subscheme such that ψ_Z is identity. Let $\Omega^*_Z(X) \subset \Omega^*(X)$ be the image of the map $\Omega^*(Z) \xrightarrow{i_*} \Omega^*(X)$. Then, ψ_Z induces a map $\psi^Z_X : \Omega^*_Z(X) \to \Omega^*_Z(X)$ such that the diagram*

$$\begin{tikzcd}
\Omega^*_Z(X) \arrow[r] \arrow[d, "\psi^Z_X"'] & \Omega^*(X) \arrow[d, "\psi_X"] \\
\Omega^*_Z(X) \arrow[r] & \Omega^*(X)
\end{tikzcd} \tag{6.2}$$

is commutative and ψ^Z_X is identity.

Proof. We consider the following diagram.

$$\begin{tikzcd}
\Omega^*(Z) \arrow[r, "\pi^*_Z"] \arrow[d, "i_*"'] & \Omega^*(E_Z/B) \arrow[r, "\cdot p_N"] \arrow[d, "i'_*"] & \Omega^*(E_Z/B) \arrow[r, "\pi_{Z*}"] \arrow[d, "i'_*"] & \Omega^*(Z) \arrow[d, "i_*"] \\
\Omega^*(X) \arrow[r, "\pi^*"'] & \Omega^*(E/B) \arrow[r, "\cdot p_N"'] & \Omega^*(E/B) \arrow[r, "\pi_*"'] & \Omega^*(X).
\end{tikzcd}$$

The first square from the left clearly commutes as π_Z is the pull-back of a smooth morphism. The second square commutes by Lemma 3.2. The third square is simply the commutativity of the push-forward maps. We conclude that the big outer square commutes. Since the top and the bottom composite horizontal arrows are ψ_Z and ψ_X respectively, we get the commutative square (6.2). Moreover, ψ^Z_X is identity because ψ_Z is so. □

Lemma 6.2. *For any scheme X, there is a finite filtration by closed subschemes*

$$\emptyset = X^{n+1} \subset X^n \subset \cdots \subset X^1 \subset X^0 = X$$

such that ψ_{U^i} is identity for each $0 \le i \le n$, where $U^i = (X^i - X^{i+1})$.

Proof. We prove this by induction on the dimension of X. If X is zero-dimensional, we can use Corollary 5.3 and the argument in the proof of Proposition 5.4 to reduce to the case when $X = \mathrm{Spec}(k)$ and $E/B = G/B$. We can apply Corollary 4.3 and [10, Theorem 7.6] to get a graded $\mathbb{L}$-algebra isomorphism

$$\phi_k : \mathbb{L} \otimes_{\mathbb{Q}} \Lambda \xrightarrow{\cong} \Omega^*(G/B). \tag{6.3}$$

Let $S_{\mathbb{L}} \xrightarrow{\eta} \Lambda_{\mathbb{L}}$ denote the quotient map. Under the above isomorphism, we get for any homogeneous element $m \in \mathbb{L}^i$,

$$\psi_k(m) = \pi_* \circ \phi_k \circ \eta \, (m \otimes p_N) = \pi_* \circ \phi_k \, (m \otimes \rho_N) = \pi_* \left(\rho_N \cdot \pi^*(m)\right) = \pi_* \left(1 \otimes \rho_N\right) \cdot m,$$

where the last equality holds by the projection formula. On the other hand, as $\mathbb{L}^0$ is the one-dimensional vector space generated by the class of $[\mathrm{Spec}(k)]$, we see that $\Omega^N(G/B) \cong \mathbb{L}^0 \otimes \Lambda_N \xrightarrow{\cong} \mathbb{Q}[1 \otimes \rho_N]$ and $\pi_*(1 \otimes p_N)$ is simply the class of $[\mathrm{Spec}(k)]$ in $\mathbb{L}^0$, which is the identity element. This proves the zero-dimensional case. If X is any smooth scheme and $Y = X \times G/B$, then it follows from Corollary 4.4 and the case of $X = \mathrm{Spec}(k)$ that ψ_X is identity.

In the general case, we can find an étale cover $f : X' \to X$ such that E is trivial over X. Hence $E/B \times_X X' \xrightarrow{\cong} E/B \times X'$ and $\pi_{X'}$ is just the projection map by Lemma 5.1. Since f is generically finite, we can find a dense open subset $U \overset{j}{\hookrightarrow} X$ such that U is smooth and the map $g : U' = f^{-1}(U) \to U$ is finite and étale. The proof of Lemma 6.1 shows that the right square in the diagram

$$\begin{tikzcd}
\Omega^*(U) \arrow[r, "g^*"] \arrow[d, "\psi_U"'] & \Omega^*(U') \arrow[r, "g_*"] \arrow[d, "\psi_{U'}"] & \Omega^*(U) \arrow[d, "\psi_U"] \\
\Omega^*(U) \arrow[r, "g^*"'] & \Omega^*(U') \arrow[r, "g_*"'] & \Omega^*(U)
\end{tikzcd} \tag{6.4}$$

commutes. Since g is étale, the similar argument shows that the left square commutes. Furthermore, we can apply Corollary 5.2 to choose the open subset U so that g^* is injective. We have shown above that $\psi_{U'}$ is identity. We conclude that ψ_U must also be identity.

Put $X^1 = (X - U)$. Then X^1 is a closed subscheme of X of dimension which is strictly less than that of X. The proof of the lemma now follows by induction. □

Proposition 6.3. *For any scheme X, the homomorphism ψ_X is an isomorphism.*

Proof. We choose a finite filtration of X as in Lemma 6.2. We have shown that $\psi_{(X-X^1)}$ is identity. Assume by induction that $\psi_{(X-X^m)}$ is an isomorphism and consider the diagram

$$\begin{tikzcd}
0 \arrow[r] & \Omega^*_{(X^m - X^{m+1})}(X - X^{m+1}) \arrow[r] \arrow[d, "\psi^{(X^m - X^{m+1})}_{(X - X^{m+1})}"'] & \Omega^*(X - X^{m+1}) \arrow[r] \arrow[d, "\psi_{(X - X^{m+1})}"] & \Omega^*(X - X^m) \arrow[r] \arrow[d, "\psi_{(X - X^m)}"] & 0 \\
0 \arrow[r] & \Omega^*_{(X^m - X^{m+1})}(X - X^{m+1}) \arrow[r] & \Omega^*(X - X^{m+1}) \arrow[r] & \Omega^*(X - X^m) \arrow[r] & 0
\end{tikzcd} \tag{6.5}$$

which is commutative by Lemma 6.1. The top and the bottom rows are exact by Theorem 2.1. The left vertical map is identity by Lemmas 6.1 and 6.2. The right vertical map is isomorphism by induction. We conclude that the middle vertical map is an isomorphism too and the induction continues. □

Corollary 6.4. *For any scheme X, the pull-back map $\Omega^*(X) \xrightarrow{\pi^*} \Omega^*(E/B)$ is injective.*

Proof. This follows directly from Proposition 6.3. ☐

Proposition 6.5. *Let $X \times G/B \xrightarrow{\pi} X$ be the trivial flag bundle. Then the map λ_X is an isomorphism.*

Proof. The surjectivity of λ_X and δ_X follows directly from Proposition 5.4 and Remark 5.5. So we only need to show the injectivity. By Remark 5.5 again, it suffices to show that ϕ_X is injective (and hence isomorphism). By Corollary 4.4, ϕ_X is same as the map of graded $\Lambda_{\mathbb{L}}$-modules

$$\Omega^*(X) \otimes_{\mathbb{Q}} \Lambda \xrightarrow{\phi_X} \Omega^*(E/B) \tag{6.6}$$

and this map factorizes the composite map θ_X as

$$\theta_X : \Omega^*(X) \otimes_{\mathbb{L}} S_{\mathbb{L}} \overset{\eta_X}{\twoheadrightarrow} \Omega^*(X) \otimes_{\mathbb{L}} \Lambda_{\mathbb{L}} \xrightarrow{\phi_X} \Omega^*(E/B). \tag{6.7}$$

It suffices to show that ϕ_X is injective on each graded piece of the left term in (6.6). So let

$$x \in (\Omega^*(X) \otimes_{\mathbb{Q}} \Lambda)^m = \overset{N}{\underset{p=0}{\oplus}} \Omega^{m-p} \otimes_{\mathbb{Q}} \Lambda_p.$$

be such that $\phi_X(x) = 0$.

Let $\{b^p_1, \cdots, b^p_{s_p}\}$ be a chosen $\mathbb{Q}$-basis of Λ_p and let $\{u^p_1, \cdots, u^p_{s_p}\}$ their homogeneous lifts in S. We have $b^0_1 = \rho_0 = 1$ and $b^N_1 = \rho_N$. Similarly, $u^0_1 = p_0 = 1$ and $u^N_1 = p_N$. We can then write x uniquely as

$$x = \sum_{p=0}^{N} x_p, \quad \text{where} \quad x_p = \sum_{i=1}^{s_p} x^p_i \otimes b^p_i.$$

We show inductively that $x_p = 0$ for each p. First of all,

$$\phi_X(x) = 0 \Rightarrow p_N \cdot \phi_X(x) = 0 \Rightarrow \theta_X(p_N \cdot x) = 0 \Rightarrow \phi_X(\rho_N \cdot x) = 0.$$

On the other hand, we have by Lemma 4.1,

$$\rho_N \cdot x_p = \sum_{i=1}^{s_p} x^p_i \otimes (b^p_i \cdot \rho_N)$$

which is zero for $p > 0$ and $x^0_1 \otimes \rho_N$ for $p = 0$. Thus we conclude that

$$\phi_X(x) = 0 \Rightarrow \phi_X(x^0_1 \otimes \rho_N) = 0 \Rightarrow \pi_* \circ \phi_X(x^0_1 \otimes \rho_N) = 0.$$

But the last equality is equivalent to saying that $\psi_X(x^0_1) = 0$ and this implies that $x^0_1 = 0$ by Proposition 6.3. In particular, $x_0 = x^0_1 \otimes b^0_1 = 0$.

We assume by induction that $x_q = 0$ for $q < p$ with $p \geq 1$ and we show that x_p is zero too. We prove this following the proof of [19] for the Chow groups. We fix an integer $1 \leq l \leq s_p$. By Lemma 4.1, there exists $c \in \Lambda_{N-p}$ such that

$$b^p_i \cdot c = \begin{cases} 0 \text{ if } i \neq l \\ \rho_N \text{ if } i = l. \end{cases}$$

In particular, we get

$$\begin{aligned} c \cdot x &= \sum_{j=p}^{N} \sum_{i=1}^{s_j} x_i^j \otimes (b_i^j \cdot c) \\ &= \sum_{i=1}^{s_p} x_i^p \otimes (b_i^p \cdot c) \\ &= x_l^p \otimes \rho_N, \end{aligned}$$

where the second equality occurs because $j + (N - p) > N$ for $j > p$. We lift c to a homogeneous element e of S. We then have as before,

$$\phi_X(x) = 0 \Rightarrow e\phi_X(x) = 0 \Rightarrow \theta_X(e \cdot x) = 0 \Rightarrow \phi_X(c \cdot x) = 0. \tag{6.8}$$

We conclude from this that $\phi_X(x) = 0 \Rightarrow \phi_X(x_l^p \otimes \rho_N) = 0$, which in turn implies that

$$\psi_X(x_l^p) = \pi_* \circ \phi_X\left(c_1(p_N) \cap \pi^*(x_l^p)\right) = \pi_* \circ \phi_X(x_l^p \otimes \rho_N) = 0.$$

It follows from Proposition 6.3 that $x_l^p = 0$ and the induction continues to show that $x_p = 0$. This completes the proof. □

Proof of Theorem 3.4: The surjectivity of λ_X follows from Proposition 5.4. Furthermore, the map δ_X is also surjective by Remark 5.5. Thus we only need to show that ϕ_X is injective.

Put $Y = E/B$. It follows from Lemma 5.1 that the pull-back $Y \times_X Y \to Y$ is the trivial flag bundle $G/B \times Y \xrightarrow{\pi'} Y$. We now consider the diagram

$$\begin{tikzcd} \Omega^*(X) \otimes_{S_{\mathbb{L}}^W} S_{\mathbb{L}} \arrow[r, "\phi_X"] \arrow[d, "\pi^*\otimes \mathrm{Id}"'] & \Omega^*(E/B) \arrow[d, "\pi'^*"] \\ \Omega^*(Y) \otimes_{S_{\mathbb{L}}^W} S_{\mathbb{L}} \arrow[r, "\phi_Y"'] & \Omega^*(G/B \times Y) \end{tikzcd} \tag{6.9}$$

which is commutative by Lemma 3.3. Since $S_{\mathbb{L}}$ is flat over $S_{\mathbb{L}}^W$, the left vertical arrow is injective by Corollary 6.4. The bottom horizontal arrow is injective by Proposition 6.5. We conclude that the top horizontal arrow is injective too. □

Corollary 6.6. *Let G be a connected linear algebraic group with a split maximal torus T. If $E \xrightarrow{p} X$ is a principal G-bundle over a scheme X, then*

$$\Omega^*(E) \xrightarrow{\cong} \Omega^*(X) \otimes_{C(G)} \mathbb{L} \cong \frac{\Omega^*(X)}{I\Omega^*(X)},$$

where I is the ideal of $C(G)$ generated by the Chern classes of G-homogeneous line bundles. This is an $\mathbb{L}$-algebra isomorphism if X is smooth.

Proof. By [10, Theorem 7.4], the natural map

$$\Omega_T^*(E) \otimes_{C(T)} \mathbb{L} \to \Omega^*(E)$$

is an isomorphism, which is an $\mathbb{L}$-algebra isomorphism if X is smooth. On the other hand, it follows from Theorem 2.2 that the term on the left is same as $\Omega^*(E/B) \otimes_{C(T)} \mathbb{L}$. The corollary now follows from Theorem 3.4. □

Corollary 6.7. *Let G be a connected algebraic group (not necessarily linear) over k and let $G \xrightarrow{\alpha_G} A(G)$ be the albanese morphism of G. Then there is an $\mathbb{L}$-algebra isomorphism*

$$\Omega^*(G) \cong \frac{\Omega^*(A(G))}{I\Omega^*(A(G))}.$$

Proof. It follows immediately from Corollary 6.6 and the fundamental exact sequence

$$1 \to G_{\mathrm{aff}} \to G \xrightarrow{\alpha_G} A(G) \to 1,$$

where G_{aff} is the largest connected linear algebraic subgroup of G. □

7. Cobordism of E/P

In this section, we complete the proof of Theorem 1.1 by deducing it from Theorem 3.4. This is done by using the following general technique, which the author learned from an unpublished note [6] of Edidin and Larsen. So let (G, T, B, W) be our given datum, where we have already reduced to the case when G is reductive. Since T is split, the group G is given by its root system $\Phi(G, T)$. Let Δ be a base of Φ and B the corresponding Borel. By the well known theory of root system (*cf.* [17]), for every subset I of Δ, there exists a corresponding parabolic subgroup $P_I \supset B$ and every parabolic subgroup containing B is of this form. Let Ψ_I be the intersection of Φ with the span of I and let $P_I = M_I N_I$ be the corresponding Levi decomposition, where M_I contains T and $\Phi(M_I, T) = \Psi_I$. The Weyl group of M_I with respect to T is the subgroup of W generated by the simple reflections corresponding to the elements of I. The Borel subgroup B_I of M_I corresponding to the positive roots in Ψ_I is the intersection of B with M_I and hence the natural map $P \to M_I$ gives an isomorphism $P_I/B \xrightarrow{\cong} M_I/B_I$. Since every Borel subgroup is conjugate to B, we can assume without loss of generality that P_I is same as P. We then obtain tower of fibrations

$$E \to E/N \to E/B \xrightarrow{g} E/P \xrightarrow{f} X. \tag{7.1}$$

We need the following Corollary of Theorem 3.4 to prove the case of parabolic flag bundles.

Corollary 7.1. *Let (G, T, B, W) be the datum as above and let $E \to X$ be a principal G-bundle. The natural map $\Omega^*(X) \to \Omega^*(E/B)^W$ is an isomorphism.*

Proof. This follows immediately from Theorem 3.4 using the fact that the trivial W-module $\mathbb{Q}$ is a projective $\mathbb{Q}[W]$-module, and $M^W = Hom_{C[W]}(\mathbb{Q}, M)$ for any $\mathbb{Q}[W]$-module M. □

To prove Theorem 1.1, we notice that $E/B \to E/P$ is a M/B-bundle, where B is a Borel in M containing T. If W_P is Weyl group of M with respect to T, we obtain

$$\Omega^*(E/P) \otimes_{C(T)^{W_P}} C(T) \xrightarrow{\cong} \Omega^*(E/B) \tag{7.2}$$

by Theorem 3.4. Taking the W_P-invariants and using Corollary 7.1, we get

$$\Omega^*(E/P) \xrightarrow{\cong} \Omega^*(E/B)^{W_P} \cong \Omega^*(X) \otimes_{C(T)^W} C(T)^{W_P}.$$

This completes the proof of Theorem 1.1. □

8. Higher Chow groups of flag-bundles

Let X be a scheme and let $p : E \to X$ be a principal G-bundle and let $\pi : E/B \to X$ be the flag bundle associated to a Borel subgroup of G. In this section, we describe the higher Chow groups of E and E/B in terms of the higher Chow groups of X and the characteristic classes of the maximal torus T. From this, we obtain formula for the higher Chow groups of the flag bundles $\pi : E/P \to X$.

Recall from [1] that the higher Chow groups of X are given by the homology groups $\mathrm{CH}^i(X, n) = \mathrm{H}_n(Z^i(X, \bullet))$ of the cycle complex $Z^i(X, \bullet)$ of codimension

i cycles in the simplicial spaces $X \times \Delta^{\bullet}$. We refer to *loc. cit.* for more detail and for some standard functorial properties. For a scheme X with G-action, the equivariant higher Chow groups $\mathrm{CH}^*_G(X, \bullet)$ are defined in [5]. We refer the reader to [8] for more details about these groups. We denote by $\mathrm{CH}^*(X)$, the full Chow groups $\overset{\infty}{\underset{n=0}{\oplus}} \mathrm{CH}^*(X, n)$ of X. This is a $\mathrm{CH}^*(k)$-algebra if X is smooth.

Let $S = \mathrm{CH}^*_T(k) \cong \mathbb{Q}[x_1, \cdots, x_r]$ and $S^W = S(G) = \mathrm{CH}^*_G(k) \cong \mathbb{Q}[\sigma_1, \cdots, \sigma_r]$ be as before. Then the same construction as in Section 3 (or as in [19]) gives a natural map of S-modules

$$\alpha_X : \mathrm{CH}^*(X) \otimes_{S^W} S \to \mathrm{CH}^*(E/B). \tag{8.1}$$

Lemma 8.1. *Let $E/B \times X \xrightarrow{\pi} X$ be the trivial flag bundle. Then α_X is an isomorphism.*

Proof. We first assume $X = \mathrm{Spec}(k)$. By [9, Theorem 1.6], the natural map $\mathrm{CH}^*_G(G/B) \otimes_{S^W} \mathbb{Q} \to \mathrm{CH}^*(G/B)$ is an isomorphism since G/B is smooth and projective. On the other hand, we have $\mathrm{CH}^*_G(G/B) \cong \mathrm{CH}^*_T(k)$ by [8, Corollary 3.2]. In particular, we get

$$\begin{aligned} \mathrm{CH}^*(k) \otimes_{S^W} S &\cong (\mathrm{CH}^*_T(k) \otimes_S \mathbb{Q}) \otimes_{S^W} S \\ &\cong \mathrm{CH}^*_T(k) \otimes_S (S \otimes_{S^W} \mathbb{Q}) \\ &\cong \mathrm{CH}^*_T(k) \otimes_{S^W} \mathbb{Q} \\ &\cong \mathrm{CH}^*(G/B), \end{aligned}$$

where the first isomorphism holds by [9, Theorem 1.6]. If $G/B \times X \xrightarrow{\pi} X$ is the trivial bundle, then we have

$$\begin{aligned} \mathrm{CH}^*(X) \otimes_{S^W} S &\cong \big(\mathrm{CH}^*(X) \otimes_{\mathrm{CH}^*(k)} \mathrm{CH}^*(k)\big) \otimes_{S^W} S \\ &\cong \mathrm{CH}^*(X) \otimes_{\mathrm{CH}^*(k)} (\mathrm{CH}^*(k) \otimes_{S^W} S) \\ &\cong \mathrm{CH}^*(X) \otimes_{\mathrm{CH}^*(k)} \mathrm{CH}^*(G/B) \\ &\cong \mathrm{CH}^*(X \times G/B), \end{aligned}$$

where the third isomorphism follows from the case of $X = \mathrm{Spec}(k)$ and the last isomorphism follows from [8, Lemma 3.6]. This completes the proof. □

Proof of Theorem 1.2: As in the case of cobordism, we can deduce the case of parabolic subgroups from the case of Borel subgroups. So we prove the result for the flag bundles of the type $E/B \to X$. We prove by induction on the dimension of X. If X is zero-dimensional, this follows from the case of $X = \mathrm{Spec}(k)$ shown above and the analogue of Corollary 5.3 for the higher Chow groups. In general, following the proof in the cobordism case, we can find a dense open subset $U \overset{j}{\hookrightarrow} X$ and a finite étale cover $g : U' \to U$ such that the pull-back of E/B is the trivial flag bundle on U' and the result holds for U' as shown above. We deduce the result for U by the higher Chow groups analogue of Corollary 5.3. Let Z be the complement of U in X with the reduced closed subscheme structure. We get the following diagram of long exact localization sequences

$$\begin{tikzcd}
\mathrm{CH}^*(U) \otimes S \arrow[r, "\partial"] \arrow[d, "\alpha_U"] & \mathrm{CH}^*(Z) \otimes S \arrow[r, "i_*"] \arrow[d, "\alpha_Z"] & \mathrm{CH}^*(X) \otimes S \arrow[r, "j^*"] \arrow[d, "\alpha_X"] & \mathrm{CH}^*(U) \otimes S \arrow[r, "\partial"] \arrow[d, "\alpha_U"] & \mathrm{CH}^*(Z) \otimes S \arrow[d, "\alpha_Z"] \\
\mathrm{CH}^*(E_U/B) \arrow[r, "\partial'"'] & \mathrm{CH}^*(E_Z/B) \arrow[r, "i'_*"'] & \mathrm{CH}^*(E/B) \arrow[r, "j'^*"'] & \mathrm{CH}^*(E_U/B) \arrow[r, "\partial'"'] & \mathrm{CH}^*(E_Z/B),
\end{tikzcd}$$

where the tensor product in the top row is over the ring S^W. In particular, this row is exact by the flatness of S over S^W. Since the dimension of Z is strictly smaller than that of X, we see that the maps α_Z are isomorphisms by induction. We have shown above that the maps α_U are also isomorphisms. We conclude that α_X is an isomorphism. ☐

Corollary 8.2. *Let G be a connected linear algebraic group with a split maximal torus T. Let $E \xrightarrow{p} X$ be a principal G-bundle over a scheme X. Then*

$$\mathrm{CH}^*(E) \xrightarrow{\cong} \mathrm{CH}^*(X) \otimes_{S(G)} \mathbb{Q} \cong \frac{\mathrm{CH}^*(X)}{I\mathrm{CH}^*(X)},$$

where I is the ideal of $S(G)$ generated by the Chern classes of G-homogeneous line bundles. This is a $\mathbb{Q}$-algebra isomorphism if X is smooth.

Proof. The proof is exactly same as the proof of Corollary 6.6. ☐

The following corollary recovers a result of Brion [2, Proposition 2.8] for the ordinary Chow groups $\mathrm{CH}^*(G, 0)$ of connected algebraic groups as a special case.

Corollary 8.3. *Let G be a connected algebraic group (not necessarily linear) over k and let $G \xrightarrow{\alpha_G} A(G)$ be the albanese morphism of G. Then there is a $\mathbb{Q}$-algebra isomorphism*

$$\mathrm{CH}^*(G) \cong \frac{\mathrm{CH}^*(A(G))}{I\mathrm{CH}^*(A(G))}.$$

Proof. The proof is exactly same as the proof of Corollary 6.7. ☐

Acknowledgments. This work was done while the author was visiting the Institute Fourier, Université de Grenoble in June, 2010. He would like to thank Michel Brion for the invitation and the local financial support during his stay.

References

[1] S. Bloch, *Algebraic cycles and higher K-theory*, Adv. in Math., **61**, (1986), no. 3, 267-304.

[2] M. Brion, *On the geometry of algebraic groups and homogeneous spaces*, MathArxiv. mathAG/0909.5014v1, (2009).

[3] M. Demazure, A. Grothendieck, *Schémas en Groupes*, Lecture Notes in Math., **153**, (1970), Springer-Verlag.

[4] D. Deshpande, *Algebraic Cobordism of Classifying Spaces*, MathArxiv, mathAG/0907.4437, (2009).

[5] D. Edidin, W. Graham, *Equivariant intersection theory*, Invent. Math., **131**, (1998), 595-634.

[6] D. Edidin, M. Larsen, *On the rational Chow ring of flag bundles*, Unpublished note.

[7] J. Fogarty, F. Kirwan, D. Mumford, *Geometric Invariant Theory*, 3rd Edition, Springer-Verlag, (1994).

[8] A. Krishna, *Riemann-Roch for equivariant K-theory*, MathArxiv, mathAG/0906.1696, (2009).

[9] A. Krishna, *Equivariant K-theory and higher Chow groups of smooth varieties*, MathArxiv, math.AG/0906.3109, (2009).

[10] A. Krishna, *Equivariant cobordism of schemes*, MathArxiv, arXiv:1006.3176, (2010).

[11] M. Levine, F. Morel, *Algebraic cobordism I*, Preprint, www.math.uiuc.edu/K-theory/0547, (2002).

[12] M. Levine, F. Morel, *Algebraic cobordism*, Springer Monographs in Mathematics. Springer, Berlin, (2007).

[13] M. Levine, *Comparison of cobordism theories*, J. Algebra, **322**, (2009), no. 9, 3291-3317.

[14] M. Levine, R. Pandharipande, *Algebraic cobordism revisited*, Invent. Math., **176**, (2009), no. 1, 63-130.

[15] H. Matsumura, *Commutative ring theory*, Cambridge studies in advanced mathematics, **8**, Cambridge university press.

[16] D. Quillen, *Elementary proofs of some results of cobordism theory using Steenrod operations*, Advances in Math., **7**, (1971), 29-56.

[17] T. Springer, *Linear algebraic groups*, Second edition, Progress in Math., **9**, (1998), Birkhauser.

[18] B. Totaro, *The Chow ring of a classifying space*, Algebraic K-theory (Seattle, WA, 1997), Proc. Sympos. Pure Math., Amer. Math. Soc., **67**, (1999), 249-281.

[19] A. Vistoli, *Characteristic classes of principal bundles in algebraic intersection theory*, Duke Math. J., **58**, no. 2, (1989), 299-315.

School of Mathematics, Tata Institute of Fundamental Research, 1 Homi Bhabha Road, Colaba, Mumbai, India

E-mail address: `amal@math.tifr.res.in`